

Case Report

Use of Autologous Fat Grafting as an Adjunct to FAMM Flap for Reconstruction of a Traumatic Lip Injury by Human Bite

Humayra Z U¹, Sharmin A², Khan MA³

Abstract:

Introduction: The lips have important functional and aesthetic roles in daily living⁽¹⁾. They are the focal point of the lower face, with several aesthetic units intricately controlled by a complex series of muscles. Restoration of the whole spectrum makes reconstruction of the lip especially challenging.

Case Presentation: A 25 years old female presented with a severe tissue defect on her upper lip and left angle of mouth to Japan East West Medical College Hospital, Dhaka. She explained that 1 day back she had been involved in domestic violence and her husband had bitten her on lip. An orofacial examination revealed as Lackmann's classification of facial bite wounds type IIIa². After a proper wound excision, lip defect was covered with a Facial Artery Musculomucosal (FAMM) flap in the first stage. Oral commissure reconstruction and scar revision was done after 2 weeks as second stage, to correct macrostomia and restoration of oral competence. Autologous fat graft was performed to ensure adequate volume enhancement of the newly reconstructed lip as the third and completion stage to achieve the final aesthetic goal and patient satisfaction.

Conclusions: Patient satisfaction is a whole spectrum which encompass both functional and aesthetic restoration. Therefore, aesthetic adjunct procedures may play important role in reconstructive surgeries specially in severe deformities.

Key word: Autologous fat grafting.

1. Dr. Zaman Ummay Humayra, Assistant Professor, Department of Surgery, Japan East West Medical College Hospital, Dhaka
2. Dr. Afrina sharmin, Assistant Professor, Department of Surgery, Z. H. Sikder Womens Medical College Hospital, Dhaka
3. Dr. Mustafa Amin Khan, Assistant Professor, Department of Plastic Surgery and Burn, National Medical College Hospital, Dhaka, Bangladesh.

Correspondence to: Dr. Zaman Ummay Humayra, Assistant Professor, Department of Surgery, Japan East West Medical College Hospital, Dhaka. phone:01716711063, email:drhumayra1996@gmail.com

Background:

The lips are the focal point of the lower face, with several aesthetic units intricately controlled by a complex series of muscles. Specially in case of human bites, the size and severity of the injuries vary, ranging from small lacerations, punctures or cuts to total avulsion and loss of relatively Large chunks of tissue therefore comprise a consequence social and psychological effects³.

Case Presentation: A 25 years old female presented with a severe tissue defect on her upper lip and left angle of mouth to Japan East West Medical College Hospital, Dhaka. She explained that 1 day back she had been involved in domestic violence and her husband had bitten her on lip.

An orofacial examination revealed as Lackmann's classification of facial bite wounds type IIIa ; defect on left lateral one-third of upper lip along with commissure. Vermilion reconstruction was performed after a proper wound excision, followed by coverage with an inferiorly based Facial Artery Musculo Mucosal (FAMM) flap in the first stage [Figure 1:A, B].

After two weeks, although the FAMM flap was able to cover the lip defect, there was macrostomia and oral incompetence to liquids. To correct the above mentioned functional deformity a second stage reconstructive surgery was planned constituting Oral commissure myoplasty and scar revision [Figure 1:B,C].



Figure 1: A. Post-traumatic upper lip defect, B. Coverage of lip defect with FAMM flap (7th POD) C. 6 months after 2 staged reconstructive surgery (Volume deficit at the left lateral third of upper lip).

Inspite of restoration of functional deformity of the lip and mouth, there was a certain degree of dissatisfaction of the patient. Because of deficiency of adequate fatty tissue below FAMM flap there was a

noticeable volume deficit at the left lateral third of the upper lip. Autologous fat graft was performed to ensure adequate volume enhancement of the newly reconstructed lip as the third and completion stage to achieve the final aesthetic goal and patient satisfaction⁴ [Figure 2: A,B].




Figure 2: A. 6 Months Post-reconstructive Volume deficit at the left lateral third of upper lip, B. 2 months after autologous fat graft.

Discussion:

Lip reconstruction is driven by restoration or preservation of function and aesthetics². The lips have both aesthetic and functional features that serve to provide recognizable individualized beauty, emotional cues, sensual interactions, speech abilities, and oral continence for nutrition. The facial artery musculo-mucosal (FAMM) flap, initially described by Pribaz et al,⁵ is a versatile flap, useful for covering of a wide variety of oro-nasal mucosal defects including lips. On the other hand, autologous fat transfer to the midface has definite long-term volume augmentation results⁷. Adipose tissue, actually, is the closest to the ideal filler because it is readily available; easily obtainable, with low donor-site morbidity; repeatable; inexpensive; versatile; and biocompatible⁸. A sequential application of both the methods ensured a satisfactory

recovery from a complex lip trauma. There is an abundance of literature supporting the efficacy of fat grafting in both aesthetic and reconstructive cases. Recent studies have shown the utility of adipose-derived stem cells in the improvement of wound healing, describing their ability to regenerate soft tissues and their remodeling capacity provided by their unique cytokine and growth factor profiles⁹. In plastic surgery, lipofilling is widely used in breast augmentation and reconstruction¹⁰⁻¹³ and in volume and contour deformities of the trunk and lower limbs¹⁴⁻¹⁶. Fat grafts have an important role in the treatment of facial hemiatrophy and lipodystrophy, in recontouring and rejuvenation of the aging face and the hands, in the treatment of depressed or altered scars. It is this regenerative capacity that is of particular interest also in chronic wound, including burns and ulcers wound therapy¹⁷.

Conclusion:

As the lips have important functional and aesthetic roles in daily living^[1], correction must be done, addressing both factors making reconstruction of the lip especially challenging. Therefore, aesthetic adjunct procedures may play an important role in reconstructive surgeries specially in complex deformities.

References:

1. Neligan PC. Strategies in lip reconstruction. *ClinPlast Surg.* 2009 Jul. 36(3):477-85. [Medline].
2. Kronic AL, Weitzul S, Taylor RS. Advanced reconstructive techniques for the lip and perioral area. *DermatolClin.* 2005 Jan. 23(1):43-53, v-vi. [Medline].
3. Shubi FM, Hamza OJ, Kalyanyama BM, Simon EN. Human bite injuries in the oro-facial region at the Muhimbili National Hospital, Tanzania. *BMC Oral Health.* 2008 Apr 30;8:12. doi: 10.1186/1472-6831-8-12..
4. Fontes T, Brandão I, Negrão R, Martins MJ, Monteiro R. Autologous fat grafting: Harvesting techniques. *Ann Med Surg (Lond).* 2018;36:212–218. Published 2018 Nov 13. doi:10.1016/j.amsu.2018.11.005.
5. J. Pribaz, W. Stephens, L. Crespo, G. Gifford. A new intraoral flap: facial artery musculomucosal (FAMM) flap. *PlastReconstrSurg,* 90 (1992), pp. 421-429 Medline.
6. Lackmann GM, Draf W, Isselstein G, Töllner U. Surgical treatment of facial dog bite injuries in children. *J Craniomaxillofac Surg.* 1992;20:81–86. doi: 10.1016/S1010-5182(05)80472-X.[PubMed][CrossRef[Google Scholar].
7. Meier JD, Glasgold RA, Glasgold MJ. Autologous Fat Grafting: Long-term Evidence of Its Efficacy in Midfacial Rejuvenation. *Arch Facial Plast Surg.* 2009;11(1):24–28. doi:10.1001/archfacial.2008.518.
8. Bellini, E., Grieco, M. P., & Raposio, E. (2017). The science behind autologous fat grafting. *Annals of medicine and surgery* (2012), 24, 65–73. doi:10.1016/j.amsu.2017.11.001.
9. A. Condé-Green, A.A. Marano, E.S. Lee, T. Reisler, L.A. Price, S.M. Milner, M.S. Granick, Fat grafting and adipose- derived regenerative cells in burn wound healing and scarring: a systematic review of the literature, *Plast. Reconstr. Surg.* 137 (2016) 302–312.
10. E. Raposio, N. Bertozzi, Autologous fat grafting

- and processing after breast reconstruction, *Surg. Chron.* 22 (2017) 66–72.
11. M. Gardani, N. Bertozzi, M.P. Grieco, M. Pesce, F. Simonacci, P.L. Santi, E. Raposio, Breast reconstruction with anatomical implants: a review of indications and techniques based on current literature, *Ann. Med. Surg.* 21 (2017)96–104.
 12. N. Bertozzi, M. Pesce, P. Santi, E. Raposio, One-stage immediate breast reconstruction: a concise review, *Biomed. Res. Int.* 2017 (2017) 6486859.
 13. N. Bertozzi, M. Pesce, P.L. Santi, E. Raposio, Oncoplastic breast surgery: comprehensive review, *Eur. Rev. Med. Pharmacol. Sci.* 21 (2017) 2572–2585..
 14. E. Delay, R. Sinna, K. Chekaroua, et al., Lipomodelling of Poland's syndrome: a new treatment of the thoracic deformity, *Aesthet. Plast. Surg.* 34 (2010) 218–225.
 15. E. Grignaffini, M.P. Grieco, N. Bertozzi, M. Gandolfi, D. Palli, F.G. Cinieri, M. Gardani, E. Raposio, Post-bariatric abdominoplasty: our experience, *Acta Biomed.* 86 (2015) 278–282.
 16. M. Grieco, E. Grignaffini, F. Simonacci, E. Raposio, Analysis of complications in post-bariatric abdominoplasty: our experience, *Plast. Surg. Int.* 2015 (2015)209173.
 17. R. Jakubietz, J.G. Grünert, D.F. Kloss, R. Meffert, K. Schmidt, M.G. Jakubietz, Aging and aesthetic ideal of the hand, *Hautarzt* 60 (2009) 220–225.
-